
気象庁海洋月報

Monthly Ocean Report

May 1999 (No. 77)



Climate and Marine Department
Japan Meteorological Agency

気象庁 気候・海洋気象部

平成１１年５月

気象庁海洋月報は、気象庁 気候·海洋気象部 海洋課で作成しています。
本月報に関する問い合わせは、

〒100-8122 東京都千代田区大手町1-3-4 気象庁 気候·海洋気象部 海洋課まで。
電 話 (03) 3212-8341 (内線 5128、5129)
F A X (03) 3211-3047
電子メール ocean@hq.kishou.go.jp

Monthly Ocean Report is prepared by the Oceanographical Division, Climate and Marine Department, Japan Meteorological Agency.

Correspondence relating to this report may be directed to :

Masaro Saiki
Director, Oceanographical Division
Climate and Marine Department, Japan Meteorological Agency
1-3-4 Ote-machi, Chiyoda-ku, Tokyo 100-8122 Japan
Facsimile: +81-3-3211-3047
e-mail : ocean@hq.kishou.go.jp

表紙は、1999年5月の太平洋の月平均海面水温偏差図です。
陰影部は、平年より高温の海域を示しています。

The figure on the front cover shows the monthly mean SST anomalies for May 1999 in the Pacific Ocean.
Areas of positive anomalies are shaded.

目　次

Table of Contents

海況の特徴（１９９９年５月）

日本近海の海況

海面水温は、オホーツク海南部から千島列島周辺で平年より１℃以上低く、北海道の南から常磐沖にかけて平年より１℃以上高かった。本州の南から南西諸島にかけては-0.5℃以下の負偏差がみられた（第4図）。

黒潮は、九州から紀伊半島にかけて接岸した後、33°Nに沿って東進し、上・中旬は八丈島付近を、下旬は八丈島の北を通った後、房総半島沖を北東に流れていた（第9図）。

親潮系の冷水の沿岸寄りの分枝は39.0°N，142.5°Eまで，沖合の分枝は39.0°N，145.0°Eまで，それぞれ達していた（第5図）。

太平洋の海況

太平洋赤道域の海面水温は、150°E～105°Wで平年より低く、155°E～125°Wでは平年より0.5℃以上低かった。一方、140°E付近と100～90°Wでは、平年より0.5℃以上高かった。

北太平洋では、30°N沿いの正偏差域は先月よりも東に広がり、150°E～140°Wで平年より0.5℃以上高く、155°E～145°Wでは+1℃以上の正偏差であった。この+0.5℃以上の正偏差域はフィリピン近海まで広がっていた。一方、40°N以北では、ほぼ全域で負偏差となっていた。また、カリフォルニア沖及び20°Nに沿った145～120°Wでは平年より１℃以上低かった（第16図）。

Highlights May 1999

Seas adjacent to Japan

SSTs were more than 1°C below normal in the southern part of the Sea of Okhotsk and around the Kuril Islands. Positive SST anomalies exceeding +1°C prevailed south of Hokkaido, off Sanriku and off Joban. Negative SST anomalies exceeding -0.5°C were found from the seas south of Honshu to those around the Nansei Islands (Fig. 4).


The Kuroshio flowed along the coasts from Kyushu to the Kii Peninsula, then eastward along 33°N. After passing near Hachijo Island in the first and second 10-days of the month, and north of Hachijo Island in the last 10-day, it flowed NE-ward off the Boso Peninsula (Fig. 9).

Coastal and off-coastal branches of the Oyashio cold water extended southward to 39.0°N, 142.5°E and 39.0°N, 145.0°E, respectively (Fig. 5).

Pacific Ocean


In the equatorial Pacific, negative SST anomalies were found from 150°E to 105°W, and exceeded -0.5°C from 155°E to 125°W. Meanwhile SSTs were more than 0.5°C above normal around 140°E and between 100°W and 90°W.

In the North Pacific, positive SST anomalies along 30°N have extended eastward during the month, and those exceeding +0.5°C prevailed from the seas around the Philippines to 30°N, 140°W, and those exceeded +1°C from 155°E to 145°W. Negative SST anomalies prevailed north of 40°N. SSTs were more than 1°C below normal off California and from 145°W to 120°W along 20°N (Fig. 16).



旬平均沿岸水温(℃)
Coastal Water Temperature

地点名 (Index Number)	水温 Temp.	偏差 Anom.
石　垣(47918)	24.3	-1.3
御前崎(47655)	16.8	-0.7
八丈島(47678)	21.6	1.8
小名浜(47598)	14.8	2.8
宮　古(47585)	8.9	0.7
江　差(47428)	10.7	-
浜　田(47755)	17.0	-




海洋気象ブイロボット
旬平均海面水温(℃)
Ocean Data Buoy SST


日本海	37.9N,134.5E	13.5
四国沖	29.0N,135.0E	21.3
東シナ海	28.2N,126.3E	23.3

第1図 北西太平洋旬平均海面水温図(上)及び偏差図(下)(1999年 5月上旬)

Fig.1 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 1st ten-day period in MAY 1999. Anomalies are computed from the JMA climatology (1961-90). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.



旬平均沿岸水温(℃) Coastal Water Temperature		
地点名 (Index Number)	水温 Temp.	偏差 Anom.
石　垣(47918)	25.2	-1.0
御前崎(47655)	18.5	-0.1
八丈島(47678)	22.7	2.0
小名浜(47598)	15.2	2.3
宮　古(47585)	10.9	1.8
江　差(47428)	11.2	-
浜　田(47755)	18.4	-




海洋気象ブイロボット
旬平均海面水温(℃)
Ocean Data Buoy SST

日本海	37.9N,134.5E	15.4
四国沖	29.0N,135.0E	22.0
東シナ海	28.2N,126.3E	25.0


第2図 北西太平洋旬平均海面水温図(上)及び偏差図(下)(1999年 5月中旬)

Fig.2 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 2nd ten-day period in MAY 1999. Anomalies are computed from the JMA climatology (1961-90). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.



旬平均沿岸水温(℃)
Coastal Water Temperature

地点名 (Index Number)	水温 Temp.	偏差 Anom.
石 垣(47918)	25.8	-0.7
御前崎(47655)	17.9	-1.7
八丈島(47678)	23.1	2.0
小名浜(47598)	16.2	2.8
宮 古(47585)	11.5	1.3
江 差(47428)	12.4	-
浜 田(47755)	19.0	-




海洋気象ブイロボット
旬平均海面水温(℃)
Ocean Data Buoy SST

地点	位置	水温
日本海	37.9N,134.5E	15.2
四国沖	29.0N,135.0E	22.6
東シナ海	28.2N,126.3E	24.7

第3図 北西太平洋旬平均海面水温図(上)及び偏差図(下)(1999年 5月下旬)


Fig.3 Ten-day mean SST (top) and anomalies (bottom) in the western North Pacific for the 3rd ten-day period in MAY 1999. Anomalies are computed from the JMA climatology (1961-90). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.



第4図 北西太平洋月平均海面水温図(上)及び偏差図(下)(1999年 5月)


Fig.4 Monthly mean SST (top) and anomalies (bottom) in the western North Pacific for MAY 1999. Anomalies are computed from the JMA climatology (1961-1990). Contour interval is 1°C and additional anomaly contours of 0.5°C are shown by broken lines. Negative anomalies are spotted.

100m Temperatures in the Seas Adjacent to Japan



第5図　日本近海表層水温図(深さ100 m) (1999年 5月)


Fig.5 Subsurface (100 m) temperatures in the seas adjacent to Japan for MAY 1999. Contour interval is 1°C. Observation points are denoted by dots.



第6図　日本近海表層水温図(深さ200 m) (1999年 5月)


Fig.6 Subsurface (200 m) temperatures in the seas adjacent to Japan for MAY 1999. Contour interval is 1°C. Observation points are denoted by dots.

400m Temperatures in the Seas Adjacent to Japan



第7図　日本近海表層水温図(深さ400 m) (1999年 5月)

Fig.7 Subsurface (400 m) temperatures in the seas adjacent to Japan for MAY 1999. Contour interval is 1°C. Observation points are denoted by dots.



第８図　北西太平洋海流図（1999 年 5 月下旬）


Fig. 8　Analysis of the sea surface current in the western North Pacific for the 3rd ten-day period in May 1999. Small arrows indicate *in situ* observations. Major currents such as the Kuroshio and the North Equatorial Current are denoted by larger open arrows.

Location of the Kuroshio axis




第9図 旬別黒潮流軸図

Fig.9 Location of the Kuroshio axis



第10図 北西太平洋力学的海面高度分布図(1999年 5月14日)


Fig.10 Sea surface dynamic heights (10cm) in the western North Pacific derived from the altimeter data of TOPEX/POSEIDON (MAY 14, 1999).
1500db surface is assumed to be the surface of no motion.



第11図 北西太平洋海流ベクトル図(1999年 5月14日)


Fig.11 Sea surface current vectors in the western North Pacific (MAY 14, 1999).
Vectors are derived from the sea surface dynamic heights from TOPEX/POSEIDON and subsurface temperature distribution assuming the geostrophic relation.


Time Series of Area Mean SST in the Western North Pacific



第12図　海域別旬平均海面水温平年偏差時系列図


Fig.12 Time series of the ten-day mean sea surface temperature anomalies (°C) in the western North Pacific for the areas shown in the right panel. Negative anomalies are shaded. Anomalies are computed from the JMA climatology (1961-90).







第12図　海域別旬平均海面水温平年偏差時系列図（つづき）

Fig.12 (continue) Time series of the ten-day mean sea surface temperature anomalies (°C) in the western North Pacific for the areas shown in the right panel. Negative anomalies are shaded. Anomalies are computed from the JMA climatology (1961-90).










第13図 沿岸水温時系列図


Fig.13 Time series of hourly coastal water temperatures (°C) measured at 7 coastal stations of the JMA shown in the right panel.

Thin lines show ten-day mean climatology.











第14図 沿岸潮位時系列図（1999年 5月）

Fig.14 Time series of hourly sea level observation (upper line) and its anomalies (lower line) at 9 stations of the JMA in the right panel.
Short term variations are removed from raw data.

NM: new moon, FQ: first quarter
FM: full moon, LQ: last quarter
T.P.: Mean sea level of Tokyo Bay (Tokyo Peil)




Time series of Sea Level



第14図 つづき


Fig.14 continue

Ocean Data Buoy Observations



第15図　　海洋気象ブイロボットによる観測要素の時系列図（日本海：21002）
上段は３時間毎の観測値を、下段は月平均値を示す。


Fig.15　　Time series of JMA buoy observation in the Japan Sea (WMO ID No. 21002).
The upper panel shows three hourly observation and the lower panel shows the monthly mean data.



第15図　海洋気象ブイロボットによる観測要素の時系列図（四国沖：21004）
上段は３時間毎の観測値を、下段は月平均値を示す。

Fig.15　Time series of JMA buoy observation south of Shikoku (WMO ID No. 21004).
The upper panel shows three hourly observation and the lower panel shows the monthly mean data.


Ocean Data Buoy Observations



第15図　海洋気象ブイロボットによる観測要素の時系列図（東シナ海：22001）
上段は３時間毎の観測値を、下段は月平均値を示す。


Fig.15　Time series of JMA buoy observation in the East China Sea (WMO ID No. 22001).
The upper panel shows three hourly observation and the lower panel shows the monthly mean data.

SST & Anomalies in the Global Ocean




第16図　全球月平均海面水温図（上）及び偏差図（下）（1999年 5月）

Fig.16 Monthly mean sea surface temperature over the global ocean (top) and anomalies (bottom) for MAY 1999. Anomalies are computed from the JMA climatology (1961-1990). Contour interval is 1°C (top) and additional anomaly contours of ±0.5°C are shown by thin lines. Negative anomalies are spotted. Maximum coverage of sea ice is shaded. Data sources for sea ice are U.S. National Ice Center (polar regions) and JMA (the Sea of Okhotsk).




第 17 図　太平洋表層水温図（深さ 100 m）（1999 年 5 月）

Fig. 17　Subsurface (100 m) temperatures in the Pacific Ocean for May 1999. Contour interval is 1 ℃. Observation points are denoted by dots.




第 18 図　太平洋表層水温図（深さ 200 m）（1999 年 5 月）

Fig. 18　Subsurface (200 m) temperatures in the Pacific Ocean for May 1999. Contour interval is 1 ℃. Observation points are denoted by dots.



第 19 図　太平洋表層水温図（深さ 400 m）（1999 年 5 月）


Fig. 19　Subsurface (400 m) temperatures in the Pacific Ocean for May 1999. Contour interval is 1 ℃. Observation points are denoted by dots.



第 20 図　太平洋ブイ流跡図（1999 年 5 月）


Fig. 20　Movements of drifting buoys in the Pacific Ocean during May 1999. Each track is for one buoy. Small circles represent the last positions for the month.

Depth-Longitude Sections of Equatorial Temperature & Anomalies




第21図　太平洋赤道に沿う表層水温（上）及び偏差（下）の深度－経度断面図（1999年 5月）

Fig.21 Depth-longitude sections of temperature (top) and anomalies (bottom) along the equator in the Pacific Ocean for MAY 1999. Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C (top) and 0.5°C (bottom). Values greater than 28°C in the top panel are shaded. In the bottom panel, values greater than 1°C are shaded and values less than -1°C are spotted. Anomalies are computed from 1987-1998 base period means.



第 22 図　太平洋月平均海面水位偏差図（1999 年 4 月）


Fig. 22　Monthly sea level anomalies for April 1999 obtained from mean sea level (1975-1995) adjusted for atmospheric pressure. This figure is prepared by the SOC of ISLP-PAC, Department of Oceanography, University of Hawaii.



第 23 図　全球月平均海面高度偏差図（1999 年 4 月）


Fig. 23　Monthly sea surface height anomalies in the global ocean derived from the altimeter data of TOPEX/POSEIDON for April 1999. Contour interval is 5 cm. Negative anomalies are spotted. Anomalies are computed from 1993-1996 base period means.


Long-term Variation of Monthly Mean SST Anomalies



第24図 海域平均海面水温偏差の長期変動

Fig.24 Long-term variations of monthly mean sea surface temperature anomalies (°C) for the areas shown in the right panel. Anomalies are computed from the JMA climatology (1961-90). Thin lines show the monthly mean and thick lines their 13-month running means.





第25図　太平洋赤道域の海面水温偏差時系列図

Fig.25 Time series of monthly mean SST anomalies (°C) in the equatorial Pacific Ocean for the areas indicated at the bottom panel. Thin lines show the monthly means and thick lines their 5-month running means. Positive anomalies (A, B, C) and negative anomalies (D) are shaded. Anomalies are computed from the JMA climatology (1961-1990).

El Nino Monitoring Indices

MONTH	REGION D 14N-EQ 130E-150E SST	REGION D Anomaly	REGION A 4N-4S 160E-150W SST	REGION A Anomaly	REGION B 4N-4S 150W-90W SST	REGION B Anomaly	REGION C EQ-10S 90W-80W SST	REGION C Anomaly	SOI	TAHITI 17-33S 149-37W	DARWIN 12-24S 130-52E
	SEA SURFACE TEMPERATURE AND ANOMALY (°C)								SOI	SEA LEVEL PRESSURE ANOMALY (hPa)	
JAN 97	28.6	0.1	28.3	0.3	24.7	-0.7	23.4	-0.9	0.3	1.7	0.9
FEB 97	28.5	0.2	28.2	0.2	25.9	-0.3	25.9	0.2	1.2	1.0	-1.9
MAR 97	28.3	-0.1	28.4	0.3	27.0	0.1	27.1	1.2	-0.6	-0.5	0.9
APR 97	29.0	0.2	29.1	0.9	27.5	0.4	26.3	1.2	-1.1	0.4	2.3
MAY 97	29.5	0.4	29.4	0.9	27.9	1.3	26.5	2.6	-2.5	-2.1	0.9
JUN 97	29.6	0.3	29.3	0.8	28.1	2.0	25.7	3.0	-2.3	-0.8	2.1
JUL 97	29.5	0.3	29.4	0.9	27.8	2.6	25.5	3.9	-0.8	-0.1	1.2
AUG 97	29.0	-0.1	29.2	0.8	27.7	3.1	24.0	3.4	-2.0	-0.9	2.2
SEP 97	29.0	-0.2	29.4	1.1	27.5	2.9	24.2	3.9	-1.4	0.1	2.7
OCT 97	29.6	0.3	29.1	0.7	27.9	3.3	24.8	4.1	-1.6	-0.2	2.7
NOV 97	29.5	0.3	29.3	0.9	28.2	3.6	25.4	3.9	-1.2	0.4	2.7
DEC 97	29.1	0.2	29.2	1.0	28.3	3.4	27.1	4.5	-0.9	-0.5	1.1
JAN 98	28.3	-0.2	29.0	1.0	28.4	3.0	28.0	3.7	-2.3	-3.6	1.4
FEB 98	28.1	-0.2	28.8	0.8	28.8	2.6	28.8	3.1	-1.8	-0.9	3.5
MAR 98	28.3	-0.1	28.6	0.5	29.0	2.1	28.8	2.9	-2.5	-2.8	2.3
APR 98	28.8	0.0	28.5	0.3	28.9	1.8	28.5	3.4	-2.3	-1.5	1.4
MAY 98	29.3	0.2	28.7	0.2	28.3	1.7	27.6	3.7	0.0	0.4	0.5
JUN 98	29.7	0.4	28.4	-0.1	26.1	0.0	24.7	2.0	1.1	0.9	-0.5
JUL 98	29.9	0.7	28.0	-0.5	25.3	0.1	23.2	1.6	1.5	1.8	-0.6
AUG 98	29.9	0.8	27.7	-0.7	24.5	-0.1	21.7	1.1	1.0	1.4	-0.2
SEP 98	30.1	0.9	27.8	-0.5	23.9	-0.7	21.0	0.7	1.0	1.4	-0.4
OCT 98	30.0	0.7	27.3	-1.1	23.8	-0.8	20.8	0.1	1.2	0.8	-1.1
NOV 98	29.9	0.7	27.2	-1.2	24.1	-0.5	21.6	0.1	1.0	0.4	-1.5
DEC 98	29.5	0.6	27.1	-1.1	23.8	-1.1	23.2	0.6	1.5	1.1	-1.5
JAN 99	29.3	0.8	26.4	-1.6	24.2	-1.2	23.9	-0.4	1.5	1.8	-1.5
FEB 99	29.1	0.8	26.7	-1.3	25.5	-0.7	25.9	0.2	0.8	1.3	-0.6
MAR 99	29.3	0.9	27.0	-1.1	26.8	-0.1	26.4	0.5	0.8	0.4	-1.3
APR 99	29.3	0.5	27.3	-0.9	26.7	-0.4	24.1	-1.0	1.8	1.3	-0.9
MAY 99	29.7	0.6	27.8	-0.7	26.4	-0.2	23.8	-0.1	0.3	1.1	1.1

第1表　エルニーニョ監視指数

Table 1 El Nino Monitoring indices


Time-Longitude Section of SST & Anomalies



第 26 図　赤道に沿った海面水温（左）及び偏差（右）の時間-経度断面図


Fig. 26　Time-longitude sections of SSTs (left) and anomalies (right) along the equator (2°N-2°S) in the global ocean. Contour interval is 1°C (left) and 0.5°C (right). Values greater than 28°C are shaded and values less than 23°C are spotted in the left panel. Dashed contours indicate negative anomalies in the right panel. Anomalies are computed from 1961-1990 base period means.

Time-Longitude Sections of Ocean Heat Content & Anomalies (Eq.)



第27図　太平洋の赤道に沿う海洋表層の貯熱量（左）及び偏差（右）の時間－経度断面図


Fig.27 Time-longitude sections of ocean heat content in the upper 260 m along the equator (left) and anomalies (right) in the Pacific Ocean. Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C (left) and 0.5°C (right). Values greater than 20°C are shaded and values less than 16°C are spotted in the left panel. Dashed contours indicate negative anomalies in the right panel. Anomalies are computed from 1987-1998 base period means.



第28図　太平洋赤道域の表層水温の時間－深度断面図

（上：西部、中：中部、下：東部）

Fig.28 Time-depth sections of equatorial subsurface temperature for the western Pacific (top, 160°E), the central Pacific (middle, 140°W) and the eastern Pacific (bottom, 110°W). Data are derived from the Ocean Data Assimilation System of JMA. Contour interval is 1°C. Values greater than 28°C are shaded.




第29-1図　北極域の海氷図（1999年 5月14日）

Fig.29-1 Sea ice chart in the Arctic on 14 May 1999.

The shaded area represents the sea ice extent. The dotted line denotes the mean ice edge averaged for the period from 1972 through 1990.

The data source is the National Ice Center (USA).




第29-2図　南極域の海氷図（1999年 5月13日）

Fig.29-2 Sea ice chart in the Antarctic on 13 May 1999.

The shaded area represents the sea ice extent. The dotted line denotes the mean ice edge averaged for the period from 1973 through 1990.


The data source is the National Ice Center (USA).



第29-3図　オホーツク海の海氷図（1999年 5月15日）

Fig.29-3 Sea ice chart in the Sea of Okhotsk on 15 May 1999.


The dotted line denotes the mean ice edge averaged for the period from 1971 through 1995.



第29-4図　オホーツク海の海氷域面積


Fig.29-4 Time-series of 5-day sea ice extent in the Sea of Okhotsk.


: current season (1998/99)
: mean extent (1971–1995)
: maximum or minimum extent (1971–1998)
: last season (1997/98)
: mean $\pm 1\sigma$ (1971–1995)



第30図　舞鶴海洋気象台清風丸によるＰＭ線に沿った水温鉛直断面図(℃). (a) 1998年7月, (b) 1998年10月, (c) 1998年11月, (d) 1999年2月, (e) 1999年4月, (f) ＰＭ線の位置図.


Fig. 30 Vertical sections of water temperature(℃) along PM line by R/V Seifu Maru of Maizuru Marine Observatory, JMA. (a) July 1998, (b) October 1998, (c) November 1998, (d) February 1999, (e) April 1999, (f) Location of PM line.






第31-1図 測点分布図(1999年 5月)

Fig.31-1 Geographical distribution of SHIP messages (SST data, top) and BATHY/TESAC messages (subsurface temperature data, bottom) for MAY 1999.





第31-2図　測点分布図(1999年 5月)

Fig.31-2 Geographical distribution of BUOY messages (buoy drift and SST data, top) and TRACKOB messages (ship drift and SST data, bottom) for MAY 1999.

気　象　庁　海　洋　月　報

Monthly Ocean Report

May 1999 (No. 77)

本月報掲載の 1999 年 5 月のプロダクトには、気象庁が観測したデータ及び GTS（全球気象通信網）を通じて入手した国際気象通報式による通報のほか、下記の機関等から提供いただいたデータを使用しています。御礼申し上げます。

防衛庁、水産庁、海上保安庁

海洋科学技術センター、（社）漁業情報サービスセンター

東京大学

水産試験場等（北海道、岩手県、宮城県、秋田県、山形県、茨城県、東京都、新潟県、石川県、福井県、愛知県、三重県、京都府、兵庫県、和歌山県、鳥取県、山口県、佐賀県、宮崎県）

（順不同）

平成１１年６月２１日発行

編集兼発行所　気　象　庁
東京都千代田区大手町１-３-４

印刷所　モリモト印刷株式会社
東京都新宿区東五軒町３-１９